



(19)

(11) Publication number: **2001261692 A**

Generated Document.

PATENT ABSTRACTS OF JAPAN

(21) Application number: **2000070880**

(51) Intl. Cl.: **C07H 15/203**

(22) Application date: **14.03.00**

(30) Priority:

(43) Date of application
publication: **26.09.01**

(84) Designated contracting
states:

(71) Applicant: **NATL INST OF ADVANCED
INDUSTRIAL SCIENCE &
TECHNOLOGY METI**

(72) Inventor: **JOHN GEORGE
MASUDA MITSUTOSHI
SHIMIZU TOSHIMI**


(74) Representative:

**(54) METHOD FOR
PRODUCING O-GLYCOSIDE
MOLECULAR AGGREGATE**


(57) Abstract:

PROBLEM TO BE SOLVED: To obtain a new reproducible molecular aggregate usable in a wide range producible from a readily obtainable raw material by a simple method.

SOLUTION: This fibrous O-glycoside molecular aggregate is produced by dissolving an O-glycoside comprising an aldose residue as a glycosyl group and a group of general formula (1) (R is a 12-18C aliphatic saturated or unsaturated straight-chain hydrocarbon group) as an aglycone in water at an elevated temperature up to a saturated concentration and then slowly cooling the aqueous solution to cause a molecular aggregation. This spherical O-glycoside molecular aggregate is produced by further heating the molecular aggregate and sphering the heated molecular aggregate. This crystal type O-glycoside molecular aggregate is produced by heating an O-glycoside comprising an aldose residue as a glycosyl group and a group of general formula (2) (R is a 12-18C aliphatic saturated or unsaturated straight-chain hydrocarbon group) as an aglycone in the absence of a solvent to cause a crystalline molecular aggregation.



一般式(1)



一般式(2)